

TABLE OF CONTENTS

	<u>Pages</u>
Introduction	i-ii
Evaluation of Permanent Disability	1-13
• Guidelines in evaluation of pain	1
• “Disability” and “Impairment”	2
• Scheduled Injuries	3
• Administrative code DWD 80.32-3 (Formerly IND 80)	4-10
• Interpolating from the guidelines	11-12
• Pre-existing disability	12-13
• Early Return to Work	13
Reprint of Average Range of Motion Charts	14-22

For additional information about worker’s compensation, contact:

**Wisconsin Worker’s Compensation Division
Department of Workforce Development**

201 East Washington Ave.
P.O. Box 7901
Madison, WI 53707-7901
(608) 266-1340
<http://dwd.wisconsin.gov/wc>

DWD is an equal opportunity employer and service provider. If you have a disability and need assistance with this information, please dial 7-1-1 for Wisconsin Relay Service. Please contact the Worker's Compensation Division at (608) 266-1340 to request information in an alternate format, including translated to another language.

HOW TO EVALUATE DISABILITY UNDER WISCONSIN'S WORKER'S COMPENSATION LAW

INTRODUCTION

The purpose of this booklet is to assist doctors in evaluating permanent disability as provided for in Wisconsin Worker's Compensation statutes and administrative codes. Worker's Compensation Law provides for prompt medical care to an injured or ill worker plus benefits to help the worker meet expenses while they are out of work.

The program emphasizes a return to work as soon as possible and rehabilitation of the worker-patient both during and after the healing period. Insured employers must report injuries to their Worker's Compensation insurance carrier within seven days of any accident which causes more than three days of disability.

A medical report verifying details of the injury or illness is made part of the record for each claim. This doctor's report is very important because it is used by self-insured employers, insurance carriers and the WC Division to determine eligibility for compensation. The Medical Report on Industrial Injury (Form WKC-16) is used to determine the amount of WC benefits paid to replace lost wages and medical expense coverage. The doctor's role also is vital in determining the amount of compensation for permanent, partial disability that may be involved.

The sooner a needed medical report is available, the sooner appropriate benefits can be paid. The WKC-16 must be filed in any case when the temporary disability exceeds three weeks or if there is permanent disability or surgery performed as a result of the injury. If the employer or insurance carrier files a narrative report in place of the WKC-16, it should list any surgery and address the issue of permanent disability. The report is due from the self insured employer or insurance carrier within 30 days of the final payment of compensation, the Division must receive an explanation if it will be delayed.

To help doctors understand worker's compensation, several documents besides this one may be ordered from the Worker's Compensation Division, P.O. Box 7901, Madison, WI 53707.

WKC-7760: **Guide for Wisconsin Doctors, Using the WKC-16B for Worker's Compensation.**

In addition, the State Medical Society of Wisconsin publishes a form to assist doctors entitled: **Attending Physician's Return to Work Recommendations Record**. This will help doctors explain special limitations such as temperature extremes, contact with skin irritants, and visual and hearing problems the patient will face at work.

EVALUATION OF PERMANENT DISABILITY

The evaluation of disability of an injured or ill worker is a necessary part of the patient's treatment, and as such is a combination of both art and science. The final rating of the patient's disability should be the medical opinion of the doctor.

Doctors may consult rating guides on relative percentage of disability and apply more than one rating formula, but in the end the final rating will be based on the doctor's personal knowledge, experience and weighing of all anatomical damages and clinical findings.

Some elements of disability, such as range of motion, can be measured with some degree of objectivity. Here there are questions requiring exercise of professional judgment. Doctors must ask: Is the demonstrated motion by the patient consistent with the nature of the injury, the x-rays, tests and other objective findings? Is the motion consistent upon repeated testing?

Other elements of disability are more subjective and less capable of being measured precisely. Pain is a good example. There is no question that pain can be disabling. Pain is not easily measured, and judgment is required. Doctors must ask:

- Are the complaints consistent with the injury?
- Has pain been consistent since the injury?
- Does it follow an anatomic pattern?
- How does pain interfere with ability to work?
- What activities bring out the complaints?
- What is the apparent severity of the pain?
- What is an appropriate disability allowance for the element of pain?

In its **Manual for Orthopedic Surgeons in Evaluating Permanent Physical Impairment**, the American Academy of Orthopedic Surgeons suggests the following guidelines in the evaluation of pain:

Grading of Pain as a Subjective Symptom

Grade I - **Mild**: When there is a firm conviction established through thorough observation and clinical tests that pain actually exists even though there may be no organic manifestations. Pain of this degree does not contribute to physical impairment.

Grade II - **Moderate:** When the examination reveals the definite evidence of a pathological state of the involved structures that would reasonably produce the degree of pain indicated to be present. This degree of pain might require treatment and could be expected to contribute in a minor degree to permanent physical impairment.

Grade III - **Severe:** When the pathological changes and clinical findings indicate that permanent physical function is limited by pain requiring treatment for relief and contributing extensively to permanent physical impairment.

Grade IV - **Very Severe:** When the pathological changes and clinical signs indicate limitation of physical function by pain to such a degree that physical impairment is nearly complete.

While pain is probably the most frequent subjective complaint, a similar reasoning process is necessary in judging other elements of disability. Examples are loss of strength, lack of endurance, loss of sensation and sensitivity to heat or cold.

It is very important for doctors to understand the distinction between “disability” in a “workplace sense” and “impairment.” In evaluating disability, this is very significant. The American Medical Association and the American Academy of Orthopedic Surgeons have recognized the distinction as follows:

- “1. Permanent disability is not a purely medical condition. A patient is ‘permanently disabled’ or ‘under a permanent disability’ when his (or her) actual or presumed ability to engage in gainful activity is reduced or absent because of ‘impairment’ and no fundamental or marked change in the future can be expected.
2. Permanent impairment is a purely medical condition. Permanent physical impairment is any anatomical or functional abnormality or loss after maximum medical rehabilitation has been achieved, which abnormality or loss the physician considers stable or non-progressive at the time evaluation is made.”

AMA GUIDE: Journal of
American Medical Assoc.

Some idea of this important difference can be gleaned from the fact that the AMA guidelines evaluate a blind person as having an 85 percent impairment of the whole person, while the Worker’s Compensation Law states the disability is 100 percent of permanent, total disability.

SCHEDULED INJURIES

The Worker's Compensation Act establishes a schedule of allowances of weeks of compensation to be paid for amputation of the extremities (arms and legs) or any parts of the extremities, plus loss of vision or hearing. Doctors should evaluate loss of use that is less **than total amputation** as a percentage of total loss. This type of injury is referred to as a SCHEDULED INJURY, and is covered by Section 102.52 of the Wisconsin Statutes.

The first section of rule DWD 80.32 states:

“(1) The disabilities set forth in this section are the minimums for the described conditions. However, findings of additional disabling elements shall result in an estimate higher than the minimum. The minimum also assumes that the member, the back, etc. was previously without disability. Appropriate reduction shall be made for any preexisting disability. Note: An example would be where, in addition to a described loss of motion, pain and circulatory disturbance further limits the use of an arm or a leg.”

At times disabilities based on loss of motion may not fall exactly within the standards established in the rule, but can be interpolated fairly easily from the standards. For example, shoulder injury with limitation of active elevation and abduction to 90 degrees represents a 20 percent loss of use at the shoulder under the rule. (See #7, “Shoulder”.) Limitation to 135 degrees is a 5 percent loss of use. If, in an actual sense, the patient's limitation was 120 degrees, the evaluation might well be a 10 percent loss of use.

The rest of the rule, (#2 through #14) follows:

WISCONSIN ADMINISTRATIVE CODE

DWD 80.32 Permanent disabilities. Minimum percentages of loss of use for amputation levels, losses of motion, sensory losses and surgical procedures. The disabilities set forth in this section are the minimums for the described conditions. However, findings of additional disabling elements shall result in an estimate higher than the minimum. The minimum also assumes that the member, the back, etc., was previously without disability. Appropriate reduction shall be made for any preexisting disability.

Note: An example would be where in addition to a described loss of motion, pain and circulatory disturbance further limits the use of an arm or a leg. A meniscectomy in a knee with less than a good result would call for an estimate higher than 5% loss of use of the leg at the knee. The same principle would apply to surgical procedures on the back. The schedule of minimum disabilities contained in this section was adopted upon the advice of a Worker's Compensation Advisory Council subcommittee after a survey of doctors experienced in treating industrial injuries.

II. *Amputations*, upper or lower extremities

At functional level	Equivalent to amputation at mid point
Stump unsuitable to accommodate prosthesis	Equivalent to amputation at next most proximal joint
Stump not functional	Grade upward

All ranges of joint motion or degrees of ankylosis not listed below are to be interpolated from existing percent of disability listed.

III *Hip*

Ankylosis, optimum position, generally 15° to 30° flexion	50%
Mal position	Grade upward
To compute disabilities for loss of motion relate % of motion lost to average range	
Shortening of leg (no posterior or lateral angulation)	
No disability for shortening less than 3/4 inch	
3/4 inch	5%
1 inch	7%
1-½ inches	14%
2 inches	22%

Greater than 2 inches of shortening results in greater proportionate rating than above

Prosthesis	Total	Minimum of 40%
	Partial	35%

IV. *Knee*

Ankylosis, optimum position, 170°	40%
Remaining range, 180° - 135°	25%
Remaining range, 180° - 90°	10%
Prosthesis	Total 50%
	Partial 45%
Removal of patella	To be based on functional impairment
Total or partial meniscectomy (open or closed procedure)	
Excellent to good result	5%
Anterior cruciate ligament repair	Minimum of 10%

V. *Ankle*

Total ankylosis, optimum position, total loss of motion	40%
Ankylosis ankle joint	
Loss of dorsi and plantar flexion	30%
Subtalar ankylosis	
Loss of inversion and eversion	15%

VI. *Toes*

Ankylosis great toe at proximal joint	50%
All other toes at proximal	40%
Ankylosis great toe at distal joint	15%
All other toes at any interphalangeal joint	If no deformity, no disability
Mal position	On merits
Loss of motion	No disability

VII. *Shoulder*

Ankylosis, optimum position, scapula free	55%
In mal position	Grade upward
Limitation of active elevation in flexion and abduction to 45° but otherwise normal	30%
Limitation of active elevation in flexion and abduction to 90° but otherwise normal	20%
Limitation of active elevation in flexion and abduction to 135° but otherwise normal	5%
Prosthesis	50%

VIII. *Elbow*

Ankylosis, optimum position, 45° angle	
With radio-ulnar motion destroyed	60%
With radio-ulnar motion intact	45%
Rotational ankylosis in neutral position	20%
Any mal position	Grade upward
Limitation of motion elbow joint, radio-ulnar motion unaffected	
Remaining range - 180° - 135°	35%
Remaining range - 135° - 90°	20%
Remaining range - 180° - 90°	10%
Rotation at elbow joint	
Neutral to full pronation	10%
Neutral to full supination	15%

IX.	<i>Wrist</i>	
	Ankylosis, optimum position 30° dorsiflexion	30%
	Mal position	Grade upward
	Total loss dorsiflexion	12 ½%
	Total loss palmarflexion	7 ½%
	Total loss inversion	5%
	Total loss eversion	5%

X.	<i>Complete Sensory Loss</i>	
	Any digit	50% Lesser involvement to be graded appropriately -35% for palmar, 15% for dorsal surface
	Total median sensory loss to hand	65-75%
	Total ulnar sensory loss to hand	25%
	Ulnar nerve paralysis	
	Above elbow, sensory involvement	50% at wrist
	Below elbow, motor and sensory involvement	45-50% at wrist
	Below elbow, motor involvement only	35-45% at wrist
	Below elbow, sensory involvement only	5-10% at wrist
	Median nerve paralysis	
	Above elbow, motor and sensory involvement	55-65% at wrist
	Thenar paralysis with sensory loss	40-50% at wrist
	Radial nerve paralysis	
	Complete loss of extension, elbow wrist and fingers	45-55% at shoulder
	Complete loss of extension, wrist and fingers	45-55% at wrist
	Peroneal nerve paralysis	
	At level below knee	25-30% at knee

XI. *Back*

Removal of disc material, no undue symptomatic complaints or any objective findings	5%
Chymopapain injection	To be rated by doctor
Spinal fusion, good results	5% minimum per level
Implantation of an artificial spinal disc	7.5%
Removal of disc material and fusion	10% per level
Cervical fusion, successful	5%
Compression fractures of vertebrae of such degree to cause permanent disability may be rated 5% and graded upward	

Note: It is the subcommittee's intention that a separate minimum 5% allowance be given for every surgical procedure (open or closed, radical or partial) that is done to relieve from the effects of a disc lesion or spinal cord pressure. Each disc treated or surgical procedure performed will qualify for a 5% rating. Due to the fact a fusion involves 2 procedures a 1) laminectomy (discectomy) and a 2) fusion procedure, 10% permanent total disability will apply when the 2 surgical procedures are done at the same time or separately.

Examples:

Patient A	12/01/1990	Laminectomy	5% PTD
	05/01/1992	Fusion	increases to 10% PTD
Patient B	12/01/1990	Laminectomy & Fusion	10% PTD
	05/01/1992	Re-fusion	increases to 15% PTD
	12/01/1992	Laminectomy at New Level	increases to 20% PTD
	05/01/1993	Fusion at 12/1/92 Level	increases to 25% PTD
	12/01/1993	Re-fusion at 5/1/93 Level	increases to 30% PTD

XII. *Fingers*

(a) *Complete ankylosis*

Thumb	Mid-position	Complete Extension
Distal joint only	25%	35%
Proximal joint only	15%	20%
Distal and proximal joints	35%	65%
Carpometacarpal joint only	20%	20%
Distal, proximal and carpometacarpal joints	85%	100%

Fingers	Mid-position	Complete Extension
Distal joint only	25%	35%
Middle joint only	75%	85%
Proximal joint only	40%	50%
Distal and middle joints	85%	100%
Distal, middle and proximal joints	100%	100%

(b) Loss of Motion

Fingers	Loss of Flexion	Loss of Use	Loss of Extension	Loss of Use
Distal joint only	10% -	1%	10% -	2%
	20% -	2%	20% -	4%
	30% -	3%	30% -	6%
	40% -	5%	40% -	8%
	50% -	10%	50% -	15%
	60% -	15%	60% -	20%
	70% -	20%	70% -	30%
	80% -	25%	80% -	40%
			100% -	60%
Middle joint only	10% -	5%	10% -	2 ½%
	20% -	10%	20% -	5%
	30% -	15%	30% -	10%
	40% -	25%	40% -	15%
	50% -	40%	50% -	30%
	60% -	50%	60% -	50%
	70% -	60%	70% -	70%
	80% -	70%	80% -	90%
			100% -	100%
Proximal joint only	10% -	5%	10% -	2 ½%
	20% -	10%	20% -	5%
	30% -	15%	30% -	15%
	40% -	20%	40% -	20%
	50% -	25%	50% -	25%
	60% -	30%	60% -	40%
	70% -	35%	70% -	75%
	80% -	40%	80% -	85%
			90% -	100%

Thumb
Distal joint same as fingers
Proximal joint 40% of the loss of use indicated for fingers

XIII. Kidney

Loss of one kidney 5%
permanent total disability.

XIV. Loss of Smell

Total loss of sense of smell
2 ½% permanent total
disability.

DWD 80.33 Permanent disabilities; fingertip amputations. In estimating permanent disability as a result of fingertip amputations, amputation of the distal one-third or less shall be considered the equivalent of 45% loss of use of the distal phalanx, amputation of not more than the distal two-thirds but more than the distal one-third shall be considered the equivalent of 80% loss of use of the distal phalanx, and amputation of more than the distal two-thirds shall be considered as 100% loss of the distal phalanx, provided there is not added disability as a result of malformed nail or tissue. In no case shall the allowance be greater than it would have been for amputation of the entire distal phalanx.

ADDITIONAL ITEMS TO ASSIST DOCTORS

1. Interpolating from the guidelines.

Having used the guidelines in the rule as starting points, doctors must then ask what other elements of disability are present and what would constitute an appropriate allowance. Questions doctors should ask are:

- Is pain present?
- How extreme is the pain?
- Does it limit activity and to what extent?
- Is it constant or intermittent?
- What is my professional judgment on an appropriate disability evaluation for pain?
- Is there altered or absent sensation?
- Is there sensitivity to heat or cold?
- Are there unstable grafts?

Ankylosis (the fused or frozen position) of the wrist in 30 degrees of dorsiflexion (bent up) is a 30 percent loss of use at the wrist. If an injury involves loss of palmar flexion (palm down), dorsiflexion and radial deviation, the question to be asked is how the disability compares with ankylosis in the “position of function.” This is the fused position which would allow the most use, or the “most favorable” position. Is it more or less? How much more or less?

In interpolating the guidelines, it is important to remember that the first few degrees of motion from the “position of function” are more important than the degrees lost at the extremes of the normal range of motion. Motion measurements are based on active, not passive, motion. Motion may be effectively limited by pain, even though greater motion is mechanically possible.

It is virtually impossible to develop guidelines that will cover, in whole or in part, every conceivable condition. In those situations not covered, doctors must rely upon professional judgment just as that professional judgment is relied upon in treatment. There is no substitute. Patients with very similar pulmonary function tests, for example, may have markedly different degrees of dyspnea (shortness of breath). A person with Raynaud’s phenomenon may have normal motion but have very limited use of the hands due to the pain produced by extended activity.

In one burn case, an attending doctor described a disability as follows:
 “I restricted ‘X’ from working in extremes of heat and cold. This was done for two reasons. His burned skin is markedly more sensitive to heat and cold than is normal skin and will be damaged by extremes of heat and cold. Secondly, his joint stiffness will be markedly exacerbated by extremes of cold....’X’ will be left with permanent residuals from his injury which will consist of altered sensations of the skin, increased rigidity of the skin, intolerance to heat and cold and hypersensitivity to sunlight.”

Obviously there is a residual industrial disability that affects the employability of this injured person. The extent of that disability must be evaluated by the doctor, relying on professional knowledge and experience. Failure to arrive at a disability rating could mean that an employee with a compensable disability is denied permanent disability benefits.

Guidelines to help doctors evaluate permanent impairment have been published by the American Medical Association and the American Academy of Orthopedic Surgery. These guides may be consulted for information and comparison purposes. They are *NOT*, however, to be the basis for evaluation; the schedules are to be used for that purpose. The AMA and AAOS guides evaluate impairment, not disability. This may not be as significant as it appears since similar physical conditions result in very comparative percentage evaluations in some instances. Examples are:

	AMA	AAOS	DWD 80.32
Total ankylosis of the ankle	40%	40%	40%
Excision of cervical disc	5%	10%	5%
Ankylosis of wrist at 30 degrees dorsi-flexion	25%	35%	30%

2. Pre-existing disability

An employer or its insurer is not liable for any pre-existing disability clearly shown to exist. Employers take responsibility for injured workers as they find them. If workers are more susceptible than normal to injury because of a pre-existing condition, the employer is nevertheless liable for the effects of the injury. An employer is liable to the extent that an injury aggravates and accelerates a pre-existing condition.

Permanent total disability (PTD) means complete disability for industrial or other employment-related purposes, although the individual may be able to perform functions not related to work, such as self care, quite satisfactorily. PTD is defined in sec. 102.44(2) of Wisconsin Statutes as:

“Total impairment for industrial use of both eyes, or the loss of both arms at or near the shoulder, or of both legs at or near the hip, of one arm at the shoulder and one leg at the hip...” The Statute then goes on to say: “This enumeration is not exclusive, but in other cases the department [DWD] shall find the facts.”

In determining what standard is to be applied “in other cases” under the statute, DWD and the Wisconsin Supreme Court have determined that:

“An employee who is so injured that he (or she) can perform no services other than those which are so limited in quality, dependability, or quantity that a reasonably stable market for them does not exist, may well be classified as totally disabled.” (**Balczewski vs. DILHR**, 76 Wis. 2d 487, 493.)

3. Early return to work.

A primary purpose of Worker’s Compensation is returning the injured or ill worker as soon as possible to productive employment, and the doctor’s evaluation plays a major role in this goal.

Performing restricted work, with limits on how many pounds can be lifted or how often bending can occur, for example, facilitates return to work at the end of a healing period. When the doctor specifies the patient’s capacities and abilities, as well as restrictions (either temporary or permanent) it benefits the injured worker, the employer and the insurer.

Capacities and restrictions need to be defined in terms of time periods (i.e., two hours per day for one week followed by... etc.). Patients need to know how such recommendations affect their activities while away from the workplace, as well.

If it is the doctor’s opinion that a patient will not be able to perform the former job permanently, this recommendation should also be explained and reported. Then the vocational rehabilitation process can be initiated. This recommendation should be reflected in the percentage of permanent disability assigned to the case.

However, if a recommendation for a permanent job change is recognized before the amount of permanent disability is assigned, the recommendation should be made immediately, before the end of the healing period, as well as when assigning permanent disability.

The final section of this booklet provides doctors with a ready reference on the average range of motion in the various joints. This information is important to properly evaluate permanent disability in work-related injuries. **DWD is indebted to the Wausau Insurance Companies for allowing the state to use their drawings. They are reprinted with permission.**

Joint Measurements

To properly evaluate permanent disabilities in industrial cases, it is necessary to know the average range of motion in the various joints. A review of the literature is confusing because of the lack of standard methods of measurement.

The average motion at a given articulation must be used as the normal, since there is considerable variation dependent upon age and individual characteristics. In industrial work, the degree of motion in the affected joint should always be compared with that of the opposite joint whenever possible.

All motions should be measured from a neutral point, and the angles measured with a goniometer or a protractor.

Each arc of the range of a circle is not of equal importance from the standpoint of function. The motion in forward elevation and abduction at the shoulder is more important than adduction and backward elevation. In the knee and elbow, the completion of flexion is not of as much importance as the first 45° of motion. In the ankle a few degrees of dorsiflexion is much less disabling than the absence of dorsiflexion. The loss of the last few degrees of motion in most joints constitutes very little, if any industrial disability.

The Jaw

HINGE MOTION — open at least one inch; closed to proper occlusion. Retrusion - 1/16 inch; protrusion - 1/4 inch. Rotation through a vertical or transverse axis - 1/4 inch from midline.

The Neck




Fig. 1
Neutral
Position

Fig. 2
Forward and backward
bending (flexion and
extension)

Fig. 3
Lateral
bending

Fig. 4
Rotation to the
left and right

NEUTRAL POSITION - HEAD UP AND CHIN IN.

1. Forward bending (flexion) - 50 to 60°
2. Backward bending (extension) - 45 to 55°
3. Lateral bending, right and left - 30 to 40°
4. Rotation, right and left - 50 to 60°

When we nod the head in assent, we are seesawing the skull on the runners of the atlas joint, like a rocking chair on a tipcart. If the head is rotated from side to side in denial, it is being turned on the pivot joint between the atlas and the axis, as on a rotating platform.

The Spine




Fig. 5
Forward bending
(flexion)

Fig. 6
Backward bending
(extension)

Fig. 7
Lateral bending
right and left

Fig. 8
Rotation,
right and left

NEUTRAL POSITION - spine in perpendicular position.

1. Forward bending (flexion) - 70 to 90°
2. Backward bending (extension) - 20 to 30°
3. Lateral bending, to right and left - 25 to 35°
4. Rotation, with the pelvis fixed, right and left - comparing angle made by the shoulders with the pelvis - 30 to 40°

Motion in the spine is hard to evaluate because of its compound nature. The cervical and lumbar regions are freely movable, while motion in the thoracic spine is quite limited. Forward bending is extremely variable, and depends upon the patient's general physique, habits of exercise, laxity or contracture of the hamstring muscles, etc. On forward bending it should be noted whether the lumbar spine flattens or reverses itself. Marked loss in forward bending in the lumbar region is important.

The Shoulder



Fig. 9
Neutral
position




Fig. 10
Abduction with
adduction with
elbow fixed




Fig. 11
Abduction with
scapula fixed —
elbow extended



Fig. 12
Abduction with
scapula free —
elbow extended




Fig. 13
Forward elevation
scapula fixed
elbow extended



Fig. 14
Forward
elevation with
scapula free,
elbow extended



Fig. 15
Backward
elevation —
elbow extended



Fig. 16
Adduction —
elbow extended



Fig. 17
External & internal
rotation with arm
in abduction,
elbow at 90°



Fig. 18
External & internal
rotation with arm
at side, elbow at 90°

NEUTRAL POSITION - arm at the side with humerus in the lateral plane of the body and elbow extended. Since the arm is pointing to 6 o'clock on a circle, the starting point is one of 180°.

1. Abduction, with elbow flexed 90° (glenohumeral motion) - arc 180 - 90°
Adduction, with elbow flexed 35 to 45° (glenohumeral motion) - arc 180 - 145 to 135°
2. Abduction with scapula fixed - elbow extended 90° (glenohumeral motion) - arc 180 - 90°
3. Abduction, with scapula free - elbow extended 180° (shoulder girdle motion) - arc 180°
4. Forward elevation (flexion), with scapula fixed 90° (glenohumeral motion) - arc 180 to 90°
5. Forward elevation (flexion), with scapula free 180° (shoulder girdle motion) - arc 180°
6. Backward elevation (extension) 45 to 55° (glenohumeral motion) - arc 180 - 135 to 125°
7. Adduction with elbow extended 35 to 45° - arc 180 - 145 to 135°
8. External and internal rotation, with arm in abduction - arc of 180°
9. External rotation, with arm at side and elbow flexed 35 to 45° - arc 180 - 145 to 135°
Internal rotation, with arm at side and elbow flexed 70 to 90° - arc of 180 - 110 to 90°
10. Circumduction - 360° (shoulder girdle motion)

Motions of the arm at the shoulder are accomplished through the scapula, as well as through the glenohumeral joint. With fixation of the arm at the side, the forearm and hand can only function in the sagittal

plane. When the scapular motion is free, the disability is much less. Complete ankylosis at the glenohumeral joint, with the arm in the optimum functional position of abduction and forward elevation, and with the scapula free, still gives a very useful range of motion.

The Elbow

Hinge motion

NEUTRAL POSITION - forearm in extension

1. Extension to 180°
2. Flexion to 45° - arc 180 to 135°
3. Hyperextension may or may not be present



Fig. 19
Neutral position – forearm in extension



Fig. 20
Flexion

With the elbow fixed in marked or complete extension, the wrist and hand are rendered proportionally useless. The carrying angle of the elbow is the angle at the long axis of the humerus and the long axis of the forearm. The average is about 15°

The Forearm



Fig. 21
Supination



Fig. 22
Pronation



Fig. 23
Supination and Pronation – arm At side, elbow Flexed at 90°

NEUTRAL POSITION - elbow flexed at 90° and forearm midway between pronation and supination (forearm at the vertical)

1. Supination, with wrist, from the vertical - 90°
Pronation, with wrist, from the vertical - 90°
2. Supination, without wrist, from the vertical - 85°
Pronation, without wrist, from the vertical - 50°

The radius may be regarded as the main bone of the forearm and wrist, just above the distal end of the radius, and on the inner side of the bone, there is an articular surface known as the sigmoid cavity. This is for articulation with the distal end of the ulna. The distal portion of the ulna also has a lateral articular facet. This joint permits the radius to rotate around the ulna through an arc of about 140°, the hand and the fibrocartilage moving with the radius.

The Wrist



Fig. 24
Neutral position




Fig. 25
Dorsiflexion and palmar
flexion (extension and
flexion)



Fig. 26
A. Radial flexion
(abduction)
B. Ulnar flexion
(adduction)



Fig. 27
Dorsiflexion




Fig. 28
Palmar flexion

NEUTRAL POSITION - Hand in line with forearm with palm facing downward

1. Radial flexion - 25 - 35°; arc 180 to 155 - 145°
2. Ulnar flexion - 35 - 45°; arc 180 to 145 - 135°
3. Dorsiflexion (extension) - 55 - 65°; arc 180 to 125 - 115°
4. Palmar flexion (flexion) - 60 - 70°; arc 180 to 120 - 110°

Abduction and adduction of the wrist occur almost entirely at the radio-carpal joint. Flexion and extension are also largely at this joint. The position of strength in the hand is one of moderate dorsiflexion. As the angle of dorsiflexion is lost, the flexion power of the fingers is diminished.

The Thumb




Fig. 29
A. Neutral position
B. Adduction (apposition)



Fig. 30
Normal flexion



Fig. 31
Abduction

NEUTRAL POSITION - thumb alongside forefinger and extended.

1. Extension to the straight position - 180°
 - A. Abduction (metacarpophalangeal joint) $50 - 60^\circ$
 - B. Adduction (metacarpophalangeal joint) $20 - 25^\circ$
2. Flexion
 - Proximal joint (metacarpophalangeal joint) $30 - 35^\circ$ - arc $180 - 150$ to 145°
 - Distal joint (interphalangeal joint) $70 - 90^\circ$ - arc $180 - 110$ to 90°
3. Abduction - 50 to 60° - arc $180 - 130^\circ$ to 120°

The trapezium-metacarpal joint is theoretically a part of the wrist, but its motions are, within certain limits, independent of the wrist. Abduction of the thumb, when limited, is perhaps best measured by taking the distance from the proximal joint of the index finger to the tip of the thumb in abduction and comparing it with the opposite side. One of the main functions of the hand is the grasping power between the thumb and the fingers. Loss of motion, particularly in the thumb, which interferes with this, is therefore of extreme importance. Ankylosis of the thumb in the grasping position is less disabling than loss of motion which does not permit grasping.

The Finger

Fig. 32
Extension



Fig. 33
Normal flexion



NEUTRAL POSITION - fingers in extension

1. Extension - 180°
2. Flexion
 - Proximal joints - 80 to 90° ; arc $180 - 100$ to 90°
 - Middle joints - 90 to 100° ; arc $180 - 90$ to 80°
 - Distal joints - 45 to 60° ; arc $180 - 135$ to 120°

In the fingers, the metacarpophalangeal joint plays an important role. Limitation of motion affects the entire finger in normal closure of the hand.

The Hip




Fig. 34
Neutral position

Fig. 35
Flexion with knee bent

Fig. 36
Flexion with knee extended



Fig. 37
Hyperextension

Fig. 38
Adduction

Fig. 39
Abduction

Fig. 40
External
Rotation

Fig. 41
Internal
Rotation




Fig. 42
Rotation in flexion




Fig. 43
Normal flexion
(flexor contracture)

NEUTRAL POSITION - hip and knee in extension, patella pointing forward

1. Flexion, with knee flexed to 45° - arc 180 to 135°
2. Flexion, with knee extended 70 to 90° - arc 180 - 110 to 90°
3. Hyperextension, 20 to 30° - arc 180 - 160 to 150°
4. Adduction, 30 to 45° - arc 180 - 150 to 135°
5. Abduction, 40 to 60° - arc 180 - 140 to 120°
6. External rotation, 45 to 60° - 180 - 135 to 120°
7. Internal rotation, 30° - arc 180 - 150°
8. Permanent flexion - This can only be determined with any degree of accuracy by having the pelvis fixed, so that the lumbar spine lies flat on the table.

Flexion at the hip, with the knee extended, varies considerably with different individuals, and will depend upon the amount of laxity or contracture of the hamstring tendons. Stiffness in the hips is compensated for throughout the lumbar spine. The greater the flexion deformity is, the greater the disability.

The Knee



Fig. 44
Neutral position




Fig. 45
Flexion - supine



Fig. 46
Flexion - prone

Hinge motion

NEUTRAL POSITION - complete extension, 180°

1. Flexion - supine position to 45° - arc 180 to 135°
2. Flexion prone position to 45° - arc 180 to 45°

The Ankle and Forefoot




Fig. 47
Dorsiflexion




Fig. 48
Plantar flexion



Fig. 49
Eversion



Fig. 50
Inversion

NEUTRAL POSITION - Outer border of the foot at 90° with leg and in neutral, as regards inversion and eversion.

1. Dorsiflexion, 15 to 25° - arc 180 - 165 to 155°
2. Plantar flexion, 30 to 45° - arc 180 - 150 to 135°
3. Eversion, 5 to 15° - arc 180 - 175 to 165°
4. Inversion, 30 to 40° - arc 180 - 150 to 140°

Complete ankylosis at the ankle still leaves considerable motion in the foot. Motion of even a few degrees dorsiflexion is much less disabling than if this is entirely lost.

The subastragal joint contributes a considerable amount of lateral motion in the use of the foot.

The calcaneal-astragaloscaphoid joint allows a twisting rotation to the foot. The astragalus is the fixed point and the anterior part of the foot is moved. In pronation the head of the astragalus rolls inward and downward in the socket of the scaphoid.

The mid-tarsal joints have as their chief movements, abduction and adduction of the anterior part of the foot.

The metatarsophalangeal joints contribute very little motion to the foot.

The Toes




Fig. 51
Dorsiflexion
(hyperextension)




Fig. 52
Plantar flexion
(flexion)

NEUTRAL POSITION - with the toes parallel to the foot.

Motions cannot be accurately measured. These motions are best described by comparing their limitations with the opposite side, also as to whether the motions are abnormally free or restricted.

The metatarsophalangeal joint is particularly important in the great toe. Stiffness in this joint, especially in dorsiflexion, is disabling insofar as forward propulsion of the foot is concerned.

(When a joint is ankylosed in flexion, the degree is important. It is best to describe such a deformity in terms of a flexor contracture or permanent flexion.)

Positions of Election

When it becomes necessary to immobilize a joint, this should be placed in the so-called position of election, unless for some reason this is contra-indicated. These positions are those which allow for the greatest function in the member if ankylosis should occur, the only exception being that of the foot at the ankle.

These positions, briefly, are as follows:

1. The thumb in the position of grasping.
2. The fingers in the position of grasping.
3. The wrist should be in 15 to 25° of dorsiflexion.
4. The forearm should be in moderate supination.
5. The elbow should be at 90° flexion in the desk-worker and 120° of extension in the laborer. If both elbows are involved, one should be placed in each position.
6. The shoulder should be put in 45° of abduction and forward elevation and in a neutral position insofar as rotation is concerned.
7. The ankle should be in 10 to 15° of dorsiflexion or at least with the foot at a right angle to the leg, and preferably with the foot in the neutral position.
8. The knee should be put in 15 to 20° of flexion.
9. The hip should be put in 20° of flexion, with 5 to 10° of abduction and in a neutral position insofar as rotation is concerned, or in 5° of external rotation.
10. In the back, the normal physiological curves should be preserved.